

## Second and third trimester abortion percentages by state, according to most recently published data

Researched by: Carole Novielli for Live Action News

### Alabama: 8.5 percent

In [2016](#), 564 out of 6,642 abortions were [reported](#) from 13-22 weeks *probable post-fertilization age*, as follows 561 (13-19 weeks) and 3 (21 weeks or greater).



#### INDUCED TERMINATIONS OF PREGNANCY, OCCURRING IN ALABAMA, 2016

Note that these tables contain data for all induced terminations of pregnancy that occurred in Alabama. The women may have lived in Alabama or some other states.

Induced Terminations of Pregnancy by Probable Postfertilization Age (PPFA)	
Total	6,642
6 or Less	3,687
7	851
8	478
9	353
10	284
11	268
12	157
13	133
14	106
15	88
16	84
17	55
18	46
19	49
20	2
21	0
22	1
23	0
24	0
25	0
26 or More	0
Not Stated	0

October 26, 2017

ADPH/CHS

Page 1 of 4

Alabama abortions by gestation 2016

### Alaska: 8.1 percent

In [2017](#), 102 out of 1,255 abortions were [committed](#) from 13-16 weeks.

#### 2017 Alaska Induced Termination of Pregnancy Statistics

Table 8a: Numbers of Induced Terminations by Weeks of Estimated Gestation, 2013-2017\*

Estimated Gestation	2013	2014	2015	2016	2017
1-4	32	17	15	16	16
5-8	863	919	873	819	768
9-12	390	400	366	354	368
13-16	86	116	75	65	102
17-20	1	44	2	2	0
21-24	0	15	1	0	0
Not Stated	78	6	2	4	1
Total	1,450	1,518	1,334	1,260	1,255

2017 Alaska Abortion by gestation

## Second and third trimester abortion percentages by state, according to most recently published data

### Arizona: at least 11 percent, (residents only)

In 2016, 13,332 abortions were reported in the state with 13,170 committed on Arizona residents. Out of that number, 1,396 abortions were committed at 13 weeks or greater: 356 (13 weeks), 469 (14-15 weeks), 251 (16-17 weeks), 181 (18-20 weeks), and 139 (21 weeks and greater). The number would likely be greater if non-resident abortions were calculated in.

Table 9. Gestational age at time of abortion Arizona residents, 2016

Gestational age	Count ( N = 13,170)	Percent
≤ 8 weeks	8,793	66.8
6 weeks	4,196	31.9
7 weeks	2,706	20.5
8 weeks	1,891	14.4
9 to 13 weeks	3,325	25.2
9 weeks	1,172	8.9
10 weeks	805	6.1
11 weeks	653	5.0
12 weeks	339	2.6
13 weeks	356	2.7
14 to 15 weeks	469	3.6
14 weeks	253	1.9
15 weeks	216	1.6
16 to 17 weeks	251	1.9
16 weeks	139	1.1
17 weeks	112	0.9
18 to 20 weeks	181	1.4
18 weeks	73	0.6
19 weeks	58	0.4
20 weeks	50	0.4
≥ 21 weeks	139	1.1
Unknown	12	NA
Total	13,170	100.0

2016 Arizona Abortion by gestation

### Arkansas: 12 percent

In 2017, 376 out of 3,249 abortions were reported between 13-20 weeks *probable post-fertilization (PPF)* age, as follows: 203 (13-16 weeks), 172 (16-19 weeks), and one at 20 weeks.

Induced Abortions by Probable Post-Fertilization (PPF) and Type of Procedure  
Arkansas Occurrences - 2017

PPF_AGE	Suction	D&E Curettage	Sharp Curettage	Intrauterine Instillation	Medical (Nonsurgical)	Other Methods	Unknown Method	Total
< 5 Wks	430	0	0	0	208	0	0	638
5 Wks	334	0	0	0	286	0	0	620
6 Wks	252	0	0	0	236	0	0	488
7 Wks	263	0	0	0	157	0	0	420
8 Wks	208	0	0	0	32	0	0	240
9 Wks	168	0	0	0	1	0	0	169
10 Wks	82	0	0	0	0	0	0	82
11 Wks	110	0	0	0	0	0	0	110
12 Wks	0	106	0	0	0	0	0	106
13 Wks	0	73	0	0	0	0	0	73
14 Wks	0	68	0	0	1	0	0	69
15 Wks	0	61	0	0	0	0	0	61
16 Wks	0	53	0	0	0	0	0	53
17 Wks	0	45	0	0	0	0	0	45
18 Wks	0	46	0	0	0	0	0	46
19 Wks	0	28	0	0	0	0	0	28
20 Wks	0	1	0	0	0	0	0	1
Total	1847	481	0	0	921	0	0	3249

Arkansas abortions by post fertilization and procedure 2017

# Second and third trimester abortion percentages by state, according to most recently published data

## California: unknown

There are [no requirements](#) by California to [report](#) abortion data, but reports on taxpayer-funded abortions via Medi-Cal for [2014](#) reveal that 4438 Dilation and Evacuation ([D+E](#)) abortions were paid for by the state. In [2014](#), Guttmacher, the former "[special affiliate](#)" of Planned Parenthood, [reported](#) 157,350 abortions in California. The data does not break these abortions down by gestation.

## Colorado: at least 10 percent

The state openly admits that [published](#) abortion numbers are "believed to be significantly underestimated." Out of the 8,873 abortions reported in [2017](#), 855 were committed at 13 weeks and up: 376 (13-15 weeks), 194 (16-21 weeks), and 285 (21 and greater). These numbers are expected to be much lower than what is taking place in the state, especially later in pregnancy.

Summary characteristics of reported induced terminations of pregnancy: Colorado occurrences, 2013-2017

Characteristic	2013		2014		2015		2016		2017	
	N	%	N	%	N	%	N	%	N	%
Total Terminations	10,198	100	10,648	100	10,114	100	8,333	100	8,873	100
Residency										
Colorado	9,127	89.9	9,437	88.7	8,930	88.7	7,324	88.3	7,837	88.7
Nonresident	1,022	10.1	1,194	11.3	1,133	11.3	972	11.7	1,000	11.3
Not Stated	50	0.5	87	0.8	51	0.5	37	0.4	36	0.4
Clinical Estimate of Gestation (Weeks)										
8 or less	6,967	71	7,376	71.3	7,288	72.4	5,687	68.4	6,224	70.2
9-10	1,248	12.7	1,226	11.8	1,195	11.9	1,308	14.1	1,218	13.7
11-12	631	6.4	691	6.5	534	5.3	489	5.9	577	6.6
13-15	496	5.1	524	5.1	579	5.3	452	5.4	376	4.2
16-20	216	2.2	266	2.5	224	2.2	206	2.5	194	2.2
21+	256	2.6	308	3	298	3	277	3.3	285	3.2
Not Stated	385	3.8	297	2.8	46	0.5	14	0.2	4	0
Type of Procedure										
Suction Curettage	5,285	51.8	5,361	50.3	4,796	47.4	3,327	40.8	3,184	36.2
Medical (Non-surgical)	4,181	41	4,502	42.3	4,444	43.9	4,134	50.6	4,774	57.3
Dilation and Evacuation	509	5	492	4.6	590	5.8	440	5.4	127	1.5
Intrauterine Instillation - Saline/Prostaglandin	+	+	+	+	+	+	+	+	+	+
Sharp Curettage	+	+	+	+	+	+	+	+	+	+
Hysterotomy/Hysterectomy	+	+	+	+	+	+	+	+	+	+
Other	223	2.2	291	2.7	281	2.8	263	3.2	247	3
Not Stated	+	+	+	+	+	+	169	2	539	6.1

Colorado abortions by gestation 2013 to 2017 underestimated

## Connecticut: unknown

## District of Columbia (DC): unknown

Abortion reporting is voluntary in D.C. and the Department of Health does not receive reports on abortions committed in private physicians' offices. In [2015](#), 1,272 abortions were reported in D.C.; however, the report does not specify by weeks of gestation.

## Delaware: 8.2 percent

In [2016](#), 179 out of 2,170 abortions were [reported](#) at or after 13 weeks as follows: 162 (13-15 weeks), 14 (16-20 weeks), and 3 (21 weeks or greater).

TABLE D-17. NUMBER AND PERCENT OF INDUCED TERMINATIONS OF PREGNANCY BY SELECTED CHARACTERISTICS, DELAWARE, 2016

Characteristic	Number	Percent	Characteristic	Number	Percent
<b>Total</b>	2,170	100.0%	<b>Marital Status</b>		
<b>Place of Residence</b>			Married	251	11.6%
Delaware	1,893	87.2%	Single	1,919	88.4%
Kent	320	14.7%	Unknown	0	0.0%
New Castle	1,303	60.0%	<b>Weeks of Gestation</b>		
Sussex	270	12.4%	<7 Weeks	551	25.4%
Maryland	142	6.5%	7 Weeks	425	19.6%
New Jersey	29	1.3%	8 Weeks	335	15.4%
Pennsylvania	91	4.2%	9-10 Weeks	463	21.3%
Other State	15	0.7%	11-15 Weeks	162	7.5%
<b>Age</b>			16-20 Weeks	14	0.6%
<15	2	0.1%	21+ Weeks	3	0.1%
15-17	75	3.5%	Unknown	1	0.0%
18-19	170	7.8%	<b>Type of Procedure</b>		
20-24	625	28.8%	Suction Curettage	1,115	51.4%
25-29	599	27.6%	Medical (Non-surgical)	1,049	48.3%
30-34	401	18.5%	Intrauterine Installation	0	0.0%
35-39	222	10.2%	Other	0	0.0%
40+	76	3.5%	Unknown	6	0.3%
Unknown	6	0.0%	<b>Previous Pregnancies</b>		
<b>Race</b>			None	591	27.2%
White	1,073	49.4%	1	405	18.7%
Black	969	44.7%	2	381	17.6%
American Indian	4	0.2%	3+	793	36.5%
Asian and Pacific Islander	60	2.8%	Unknown	0	0.0%
Other Races	64	2.9%			
Unknown	0	0.0%			

2016 Delaware Abortion by gestation

## Second and third trimester abortion percentages by state, according to most recently published data

### Florida: 6.4 percent

In [2017](#), there were 68,935 elective abortions [reported](#) in the state and 4,438 were [recorded](#) in the second trimester, with one in the third trimester.

#### Agency for Health Care Administration Reported Induced Terminations of Pregnancy (ITOP) by Reason, by Trimester

2017 -- Year to Date

Reason	Total	1st Trimester	2nd Trimester	3rd Trimester
Elective Abortion	55247	52192	3055	0
Abortion Performed due to Social or Economic Reasons	11222	10456	766	0
Abortion Performed due to Emotional/Psychological Health of the Mother	911	836	75	0
Abortion Performed due to Physical Health of Mother that is not Life Endangering	700	638	62	0
Abortion Performed due to a Life Endangering Physical Condition	141	111	29	1
Abortion Performed due to Serious Fetal Genetic Defect, Deformity or Abnormality	623	201	422	0
Abortion Performed due to Incest	5	4	1	0
Abortion Performed due to Rape	86	58	28	0
<b>Florida Totals:</b>	<b>68935</b>	<b>64496</b>	<b>4438</b>	<b>1</b>

2017 Florida Abortion by gestation and reason

### Georgia: unknown

In [2016](#), Georgia reported 29,551 abortions but did not break them down further.

### Hawaii: unknown

### Idaho: 8 percent

In [2016](#), 99 out of 1,289 abortions were committed on or after the 13th week with one abortion reported in the third trimester: 97 (13-15 weeks), 1 (16-20 weeks), and 1 (25 weeks or greater).

Table of Contents

**INDUCED ABORTIONS OCCURRING IN IDAHO**  
**Clinical Estimated Weeks of Gestation and Primary Termination Procedure**  
**2006-2016**

PROCEDURE CHARACTERISTICS	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
TOTAL IN IDAHO	1,249	1,442	1,481	1,650	1,510	1,440	1,458	1,375	1,353	1,272	1,289
Weeks of gestation											
Number <sup>1</sup>											
<9	804	926	997	1,056	994	887	978	921	923	832	824
9-10	258	256	275	309	282	300	242	234	238	248	257
11-12	135	187	135	199	163	174	141	144	115	104	108
13-15	37	53	62	65	59	69	86	64	67	80	97
16-20	8	11	8	14	6	6	10	9	6	8	1
21-24	3	5	3	5	5	2	1	3	1	-	-
25+	2	-	-	1	1	-	-	-	-	-	1
Not stated	2	4	1	1	-	2	-	-	3	-	1
Termination procedure											
Number <sup>2</sup>											
Suction curettage	1,035	1,129	1,043	1,068	995	932	883	845	778	666	622
Sharp curettage (D&C)	3	2	4	5	2	2	2	3	1	-	-
Dilation and evacuation (D&E)	7	12	7	13	6	4	12	8	24	45	85
Intra-uterine instillation (saline or prostaglandin)	3	2	1	1	1	1	2	2	2	-	-
Medical (nonsurgical)	198	293	423	556	504	499	559	513	547	561	581
Hysterotomy/hysterectomy	-	-	-	1	-	-	-	-	-	-	-
Other	3	4	3	6	2	2	-	4	1	-	1
Not stated	-	-	-	-	-	-	-	-	-	-	-

Idaho abortions by gestation/procedures 2006 to 2016

## Second and third trimester abortion percentages by state, according to most recently published data

### Illinois: 11 percent (estimated)

Illinois does not segregate out the 13th week and the state reported 3,216 abortions not specified by gestation. Because this unknown number was so high, we based our calculations on all reported abortions in the 12-15 week category. In [2016](#), 4,156 abortions out of 38,382 were committed at 12 weeks or higher as follows: 2,868 (12-15 weeks), 1,154 (16-19 weeks), 134 (20-23 weeks), and 3,216 (unknown.)

2016

County Code	County Name	# per County	Age 0-19	Age 20-24	Age 25-29	Age 30-34	Age 35-39	Age 40-44	Age 45+	Age Unknown	Age 0-14	Age 15-17	Age 18-19	Married Yes	Married No	Married Unknown	Gest. 0-3	Gest. 4-7	Gest. 8-11	Gest. 12-15
600	Chicago(Zip)	15138	1348	4631	4447	2711	1519	398	*****	*****	51	461	836	958	10714	3466	*****	6959	3866	1286
000	Illinois	32663	3111	10007	9229	5762	3341	1021	97	95	104	993	2014	3267	24293	5103	*****	16029	9230	2868

\*\*\*\*\* denotes less than or equal to 50

#### denotes greater than 50 but suppressed to prevent calculation of other suppressed numbers less than or equal to 50 (\*\*\*\*\*)

2016

County Code	County Name	Gest. 16-19	Gest. 20-23	Gest. 24+	Gest. Unknown	Live Birth Yes	Live Birth No	Live Birth Unknown	Spontan. Yes	Spontan. No	Spontan. Unknown	Induced Yes	Induced No	Induced Unknown	Reason Req	Reason Other
600	Chicago(Zip)	648	67	*****	2290	9405	####	*****	####	12643	*****	####	7707	*****	14233	*****
000	Illinois	1154	134	*****	3216	20238	####	*****	####	27202	*****	####	18292	*****	31083	108

\*\*\*\*\* denotes less than or equal to 50

#### denotes greater than 50 but suppressed to prevent calculation of other suppressed numbers less than or equal to 50 (\*\*\*\*\*)

2016 Illinois Abortion Stats by gestation

### Indiana: estimated <1 percent, most likely higher

Indiana does not show how many abortions are committed at the 13th week. The state reports that in [2017](#), 1,628 abortions out of 7,778 were committed between 9-13 weeks and 28 abortions between 14-20 weeks. The percentage would likely be larger if the number of abortions at 13 weeks were specified.

In Indiana, 78.71% of terminations occurred prior to the ninth week of pregnancy, and 20.93% occurred during the 9-13 week timeframe. Terminations occurring later than 13 weeks accounted for less than 1 percent of the total for 2017.

Table 16. Estimated gestational age at time of termination in Indiana, 2017		
Weeks of Gestation	Count	Percent
≤ 8	6,122	78.71
9-13	1,628	20.93
14-15	4	0.05
16-17	8	0.10
18-20	16	0.21
<b>Total</b>	<b>7,778</b>	<b>100.0</b>

2017 Indiana Abortion by gestation



## Second and third trimester abortion percentages by state, according to most recently published data

### Iowa: 6 percent, most likely higher

Iowa does not break down abortions at 13 weeks, nor does it specify which are committed in the third trimester. In [2016](#), 3,492 abortions out of 3,722 induced abortions were committed from 9-13 weeks while 222 abortions were reported between 14-28 weeks. The percentage would likely be larger if the number of abortions at 13 weeks were specified.

TABLE 51  
GESTATIONAL AGE IN WEEKS OF FETUS BY TERMINATION TYPE  
2015-2016 OCCURRENCE DATA

Gestational Age	2016				2015			
	Surgically Induced	Medically Induced	Other Induced	Spontaneous	Surgically Induced	Medically Induced	Other	Spontaneous
Total	1,468	2,251	3	538	1,801	2,179	9	488
0 to 13 Weeks	1,247	2,242	3	508	1,546	2,177	7	457
14 to 28 Weeks	220	2	0	26	254	0	2	27
29+ Weeks	0	0	0	1	0	0	0	0
Unknown	1	7	0	3	1	2	0	4

Iowa abortions by gestation 2015 to 2016

### Kansas: 11 percent

According to a [preliminary report](#) of [reported abortions](#) in 2017, 714 out of 6,782 abortions were performed at 13 weeks and higher as follows: 415 (13-16 weeks), 289 (17-21 weeks), and 10 (22 weeks and over). Only 26 abortions were reported for physical, mental, or emotional abuse or neglect.

Selected Induced Abortion Statistics Kansas, 2017		
Selected Statistics	Number	Percent
<b>Marital Status of Patient:</b>		
Married	1,050	15.5
Unmarried	5,683	83.8
n.s.	49	0.7
Total Reported	6,782	100
<b>Weeks Gestation:</b>		
Less than 9 weeks	4,615	68.0
9-12 weeks	1,463	21.4
13-16 weeks	415	6.1
17-21 weeks	289	4.3
22 weeks & over **	10	0.1
n.s.	0	n/a
Total Reported	6,782	100
<b>Method of Abortion:</b>		
Suction curettage	2,220	32.7
Sharp curettage	12	0.2
Dilation & Evacuation	592	8.7
Medical Procedure I (Mifepristone)	3,939	58.1
Medical Procedure II (Methotrexate)	1	0.0
Intra-uterine prostaglandin instillation	0	0.0
Hysterotomy	2	0.0
Hysterectomy	0	0.0
Digoxin/Induction	0	0.0
Partial Birth Procedure**	0	0.0
Other	16	0.2
n.s.	0	n/a
Total Reported	6,782	100
<b>Reason for Determining Gestational Age (method):</b>		
Ultrasound	6,745	99.5
Ultrasound and last menstrual period	3	0.0
Ultrasound and medical records from Physician	1	0.0
Date of last menstrual period and termination date	22	0.3
n.s.	11	0.2
Total Reported	6,782	100
<b>Report of Physical, Mental, or Emotional Abuse or Neglect Filed</b>		
No	6,756	99.6
Yes	26	0.4
n.s.	0	n/a
Total Reported	6,782	100

\*\* There were 10 reported abortions performed by providers out of state to Kansas residents.

\*\* No Partial Birth procedures have been reported in Kansas since October 1999.

2017 Kansas Abortion by gestation

## Second and third trimester abortion percentages by state, according to most recently published data

Kentucky: unknown

**Louisiana: 9 percent** In [2017](#), 788 out of 8,706 abortions were committed at 13 weeks and greater: 707 (13-16 weeks) and 81 (17-19 weeks).

INDUCED TERMINATIONS OF PREGNANCY BY WEEKS OF GESTATION, RACE, AGE, AND MARITAL STATUS REPORTED OCCURRING IN LOUISIANA, 2017																	
WEEKS OF GESTATION	RACE	TOTAL	AGE										MARITAL STATUS				
			UNDER 15	15-19	20-24	25-29	30-34	35-39	40-44	45 & OVER	UNKNOWN	MARRIED	UNMARRIED	UNKNOWN			
All	All	8,706	27	791	2,592	2,532	1,631	869	250	13	<5	976	7,329	401			
	WHITE	2,623	<5	232	753	716	508	321	86	<5	-	435	2,125	63			
	NONWHITE	5,973	22	552	1,820	1,773	1,107	535	153	10	<5	520	5,138	315			
	UNKNOWN	110	<5	7	19	43	16	13	11	-	-	21	66	23			
8 AND UNDER	All	6,026	13	447	1,731	1,802	1,193	630	198	11	<5	745	4,999	282			
	WHITE	1,590	<5	158	548	543	388	221	66	<5	-	333	1,550	47			
	NONWHITE	4,025	9	285	1,175	1,230	795	399	123	8	<5	394	3,408	223			
	UNKNOWN	71	<5	<5	8	29	10	10	9	-	-	18	41	12			
9	All	731	<5	89	215	158	135	75	15	<5	-	63	637	31			
	WHITE	188	-	23	49	33	31	<5	<5	-	-	33	151	<5			
	NONWHITE	530	<5	65	162	145	100	43	11	<5	-	30	477	23			
	UNKNOWN	13	-	<5	<5	<5	<5	<5	<5	-	-	-	9	<5			
10	All	493	<5	61	159	133	79	42	16	-	-	40	419	34			
	WHITE	134	-	12	38	30	28	19	7	-	-	19	107	8			
	NONWHITE	354	<5	48	118	103	51	23	8	-	-	20	309	25			
	UNKNOWN	5	-	<5	<5	-	-	-	<5	-	-	<5	<5	<5			
11	All	379	<5	54	118	112	57	31	5	<5	-	33	332	14			
	WHITE	102	-	17	31	25	13	13	<5	-	-	9	92	<5			
	NONWHITE	273	<5	36	87	85	44	17	<5	<5	-	24	239	10			
	UNKNOWN	<5	-	<5	<5	-	<5	<5	-	-	-	<5	<5	<5			
12	All	270	<5	34	74	72	40	21	5	-	-	28	233	9			
	WHITE	63	-	5	21	16	10	8	<5	-	-	11	51	<5			
	NONWHITE	200	<5	29	74	52	29	13	<5	-	-	17	175	8			
	UNKNOWN	7	-	-	<5	<5	<5	<5	-	-	-	-	7	-			
13	All	214	<5	5	24	87	67	37	15	<5	-	23	203	8			
	WHITE	69	-	<5	28	13	18	7	<5	-	-	15	54	-			
	NONWHITE	162	<5	22	58	53	18	8	<5	-	-	8	146	8			
	UNKNOWN	<5	-	-	-	-	-	-	-	-	-	<5	<5	<5			
14	All	188	<5	27	65	46	30	16	<5	-	-	13	166	9			
	WHITE	36	<5	<5	9	9	5	9	<5	-	-	<5	33	-			
	NONWHITE	151	<5	25	56	37	24	7	<5	-	-	10	133	8			
	UNKNOWN	<5	-	-	-	-	-	-	-	-	-	-	-	<5			
WEEKS OF GESTATION	RACE	TOTAL	AGE										MARITAL STATUS				
			UNDER 15	15-19	20-24	25-29	30-34	35-39	40-44	45 & OVER	UNKNOWN	MARRIED	UNMARRIED	UNKNOWN			
15	All	161	<5	22	53	45	23	16	<5	-	-	12	143	6			
	WHITE	40	-	5	16	13	6	5	-	-	-	<5	40	<5			
	NONWHITE	113	-	17	37	30	16	11	<5	-	-	6	103	<5			
	UNKNOWN	<5	-	-	-	<5	<5	-	-	-	-	<5	<5	<5			
16	All	124	<5	14	40	25	25	<5	<5	-	-	7	113	<5			
	WHITE	29	-	7	6	5	5	<5	<5	-	-	<5	26	<5			
	NONWHITE	95	-	19	32	19	20	5	-	-	-	5	86	<5			
	UNKNOWN	<5	-	-	<5	<5	-	-	-	-	-	-	<5	<5			
17	All	62	<5	<5	11	13	7	-	-	-	-	5	35	<5			
	WHITE	9	-	-	<5	<5	<5	<5	-	-	-	<5	7	<5			
	NONWHITE	32	<5	<5	8	8	7	5	-	-	-	<5	27	<5			
	UNKNOWN	17	<5	<5	7	5	<5	<5	<5	-	-	<5	14	<5			
18	All	5	-	-	<5	<5	<5	<5	-	-	-	<5	<5	<5			
	WHITE	5	-	-	<5	<5	<5	<5	-	-	-	<5	<5	<5			
	NONWHITE	12	<5	<5	<5	<5	<5	<5	-	-	-	<5	11	-			
	UNKNOWN	-	-	-	-	-	-	-	-	-	-	-	-	-			
19	All	22	<5	<5	<5	8	<5	<5	<5	<5	-	<5	19	<5			
	WHITE	7	-	<5	<5	<5	<5	<5	-	-	-	<5	6	-			
	NONWHITE	14	<5	<5	<5	<5	<5	<5	-	-	-	<5	13	-			
	UNKNOWN	<5	-	-	-	-	-	<5	<5	<5	-	-	-	<5			
UNKNOWN	All	29	-	<5	6	6	<5	<5	<5	-	-	<5	16	<5			
	WHITE	7	-	<5	<5	<5	<5	<5	-	-	-	<5	5	-			
	NONWHITE	12	-	-	<5	<5	<5	<5	<5	-	-	<5	11	-			
	UNKNOWN	-	-	-	-	-	-	-	-	-	-	-	-	-			

## Second and third trimester abortion percentages by state, according to most recently published data

### Michigan: 13 percent

In [2017](#), 3,461 abortions out of 26,594 abortions were [committed](#) at 13 weeks and greater, including 9 abortions in the third trimester: 2,115 (13-16 weeks), 912 (17-20 weeks), 425 (21-24 weeks), 6 (25-28 weeks), and 3 (over 28 weeks).

[Home Pages](#) + [Health Statistics](#) + [Abortion Statistics](#) + [Previous](#) [Next](#)

**Table 17**  
Number and Percent of Reported Induced Abortions by Weeks of Gestation and Principal Procedure, Michigan Occurrences, 2017

Weeks of Gestation <sup>1</sup>	Principal Procedure														
	Total	Suction Curettage		Medical (Non-Surgical)		Dilation & Evacuation		Saline or Prostaglandin		Sharp Curettage (D & C)		Hysterectomy or Hysterotomy		Other Procedures	
		Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
Total	26,594	15,314	57.6	9,422	35.4	1,777	6.7	3	0.0	-	-	2	0.0	24	0.1
< 5	323	106	32.8	215	66.6	-	-	-	-	-	-	-	-	-	-
5 - 8	16,618	8,544	51.4	8,017	48.2	16	0.1	1	0.0	-	-	1	0.3	11	0.1
9 - 12	6,172	5,014	81.2	1,123	18.2	12	0.2	-	-	-	-	-	-	7	0.1
13 - 16	2,115	1,389	65.7	21	1.0	698	33.0	-	-	-	-	1	0.3	1	0.0
17 - 20	912	200	21.9	14	1.5	694	76.1	2	0.2	-	-	-	-	1	0.1
21 - 24	425	44	10.4	24	5.6	353	83.1	-	-	-	-	-	-	4	0.9
25 - 28	6	3	50.0	3	50.0	-	-	-	-	-	-	-	-	-	-
Over 28	3	3	100.0	-	-	-	-	-	-	-	-	-	-	-	-

Michigan abortion stats by gestation 2017

### Minnesota: 11.4 percent

In [2017](#), 1,159 out of 10,177 abortions were [committed](#) between 13-24 weeks including 2 in the third trimester: 557 (13-15 weeks), 450 (16-20 weeks), 150 (21-24 weeks), and 2 (25-30 weeks).

**Table 11. Clinical Estimate of Fetal Gestational Age, 2017**

	Occurring in Minnesota	Minnesota Residents
<9 weeks	7,043	6,452
9 - 10 weeks	1,241	1,123
11 - 12 weeks	654	588
13 - 15 weeks	557	492
16 - 20 weeks	450	386
21 - 24 weeks	150	122
25 - 30 weeks	2	2
31 - 36 weeks	0	0
37 weeks & over	0	0
Not Reported	80	53
Total	10,177	9,218

Minnesota abortion stats by gestation 2017



## Second and third trimester abortion percentages by state, according to most recently published data

### Mississippi: 11 percent

Although there is only one abortion facility left in the state, a report of induced abortions sent to Live Action News by the Mississippi State Department of Health shows that in 2016, 279 out of 2,569 abortions were reported at 13 weeks or greater: 272 (13-16 weeks), 3 (17-20 weeks), and 4 (21 weeks or greater).

Table F3. Reported Induced Terminations of Pregnancy and Percentage Distribution by Medical Characteristics and Race Procedures Performed in Mississippi, 2016

Medical Characteristics	Number				Percent		
	Total <sup>1</sup>	White	Black	Unknown	Total	White	Black
<b>Total</b>	<b>2,569</b>	<b>463</b>	<b>1,983</b>	<b>76</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
<b>Completed Weeks of Gestation<sup>2</sup></b>							
Under 9	1,659	319	1,253	56	64.6	68.9	63.2
9, 10	373	67	296	3	14.5	14.5	14.9
11, 12	258	42	203	7	10.0	9.1	10.2
13-16	272	33	226	10	10.6	7.1	11.4
17-20	3	1	2	0	0.1	0.2	0.1
21 or more	4	1	3	0	0.2	0.2	0.2
Unknown	0	0	0	0	-	-	-
<b>Primary Procedure</b>							
Suction Curettage	1,007	126	831	31	39.2	27.2	41.9
Medical (Nonsurgical)	1,557	335	1,149	45	60.6	72.4	57.9
Dilation and Evacuation	2	0	2	0	0.1	-	0.1
Intrauterine Instillation	0	0	0	0	-	-	-
Sharp Curettage	3	2	1	0	0.1	0.4	0.1
Hysterotomy or Hysterectomy	0	0	0	0	-	-	-
Other	0	0	0	0	-	-	-
Unknown	0	0	0	0	-	-	-

<sup>1</sup> Total includes races other than white and black

<sup>2</sup> Physician's estimate of gestational age

\*\* Less than 0.06 percent.

Mississippi abortions by gestation and procedure 2016

### Missouri: 17.3 percent

In 2016, 789 abortions out of 4,562 were committed at 13 weeks or greater as follows: 259 (13-14 weeks), 150 (15-16 weeks), 214 (17-19 weeks), 57 (20 weeks), and 109 (21 weeks or greater).

Table 12B. Recorded Abortions by Race, Age, and Type of Procedure by Weeks of Gestation: Missouri, 2016

	Total	Weeks of Gestation								
		Under 9	9-10	11-12	13-14	15-16	17-19	20	21 and Over	Unknown
<b>State Total</b>	4,562	2,224	1,015	534	259	150	214	57	109	0
<b>Patient's Race</b>										
White, Non-Hispanic	2,127	1,144	439	219	108	71	78	13	55	0
Black, Non-Hispanic	1,942	803	464	277	134	67	112	40	45	0
Other, Non-Hispanic	292	175	64	20	11	6	12	1	3	0
Unknown, Non-Hispanic	73	32	20	7	4	2	4	2	2	0
Hispanic, Any Race	128	70	28	11	2	4	8	1	4	0
<b>Patient's Age</b>										
Under 15	8	0	3	0	2	1	0	0	2	0
15	18	6	3	4	1	1	2	0	1	0
16	41	19	7	6	6	2	1	0	0	0
17	66	25	17	12	2	6	4	0	0	0
18	145	63	35	20	11	2	6	6	2	0
19	181	90	31	20	18	8	11	1	2	0
20-24	1,475	675	318	198	93	56	80	22	33	0
25-29	1,339	643	325	158	73	42	54	13	31	0
30-34	731	364	179	73	28	20	31	10	26	0
35-39	438	263	81	33	22	9	17	4	9	0
40 and Over	120	76	16	10	3	3	8	1	3	0
Unknown	0	0	0	0	0	0	0	0	0	0
<b>Type of Procedure</b>										
Curettage	2,707	1,247	775	534	148	2	1	0	0	0
Intrauterine Instillation	1	0	0	0	0	0	1	0	0	0
Hysterotomy/Hysterectomy	1	0	0	0	0	1	0	0	0	0
Laminaria D & E	626	1	0	0	111	147	205	56	106	0
Medical/Nonsurgical	1,226	976	239	0	0	0	7	1	3	0
Unknown	1	0	1	0	0	0	0	0	0	0
Other	0	0	0	0	0	0	0	0	0	0

Missouri abortion stats by gestation 2016

## Second and third trimester abortion percentages by state, according to most recently published data

**Montana: unknown**

**Nebraska: 7 percent**

In [2017](#), 133 abortions out of 1,958 were estimated between 13-21 weeks: 104 (13-16 weeks), 37 (16-20 weeks) and 1 (21 weeks and up). **The report states that there were *no* abortions determined to be medically necessary.**

**Table 19: Number and Percent of Abortions by Clinical Estimate of Gestation (in weeks) Nebraska, 2017**

<u>Estimated Gestation</u>	<u>Number</u>	<u>Percent</u>
Under 3	33	1.7
3	81	4.1
4	250	12.8
5	243	12.4
6	246	12.6
7	263	13.4
8	236	12.1
9	175	8.9
10	118	6.0
11	89	4.5
12	89	4.5
13	59	3.0
14	24	1.2
15	12	0.6
16	9	0.5
17	10	0.5
18	9	0.5
19	9	0.5
20	0	-
21	1	0.1
Not Reported	2	0.1

10

Nebraska abortion stats by gestation 2017

**Nevada: unknown**


**New Hampshire: unknown**

**New Jersey: unknown**

**New Mexico: at least 8 percent, most likely higher**

In [2016](#), the state did not break down abortions at 13 weeks, therefore the percentage is most likely greater than calculated here. 246 abortions out of 3,233 were committed at 14 weeks or greater: 508 (9-13 weeks), 94 (14-15 weeks), 55 (16-17 weeks), 42 (18-20 weeks), and 55 (21 weeks or greater). In addition, more than 272 abortions were reported as “unknown” gestation.

**Figure A-1. Percentage of Induced Abortions by Week of Gestation New Mexico Occurrence, 2016 (New Mexico Residents Only)**



New Mexico residents obtaining terminations in other states are not included in the New Mexico resident data.  
Gestation is physician's estimate.

New Mexico abortion stats by gestation 2016

# Second and third trimester abortion percentages by state, according to most recently published data

## New York: 8.1 percent

This state does not break down abortions by gestation, however they do [publish](#) abortion data by procedure and resident county. In 2015, 7,064 out of the 86,627 abortions were potentially later term procedures: 6,985 D+E, 51 [saline abortion](#), and 28 Prostaglandin.



Services News Government Local

L

Department of Health

Individuals/Families

Providers/Professionals

Health Facilities

Search

You are Here: [Home Page](#) > [Vital Statistics of New York State 2015](#) > Table 25: Induced Abortion by Operative Procedure and Resident County - New York State 2015

**Table 25: Induced Abortion by Operative Procedure and Resident County - New York State 2015**

County	Operative Procedure								
	Total <sup>1</sup>	D&C <sup>2</sup>	S&C <sup>3</sup>	D&E <sup>4</sup>	Saline Inject.	Prostaglandin	Medical	Other	Not Stated
<b>New York State</b>									
New York State	86,627	2,712	57,756	6,985	51	28	18,540	205	897
<b>New York City</b>									
New York City	58,627	1,255	42,447	4,304	32	0	10,396	14	191
Bronx	14,992	261	10,936	1,132	10	0	2,604	6	44
Kings	18,026	266	13,573	1,126	7	0	3,037	2	17
New York	10,294	159	7,625	813	5	0	1,674	0	18
Queens	13,681	559	9,053	1,111	6	0	2,844	6	111
Richmond	1,634	10	1,260	122	4	0	237	0	1
<b>Rest of State</b>									

New York State abortion stats by procedure 2015

## New York City: 11.4 percent

In [2015](#), 7,286 out 63,646 abortions reported were committed at or above 13 weeks: 3,035 (13-15 weeks), 2,766 (16-20 weeks), and 1,485 (21 weeks and greater).

## PREGNANCY OUTCOMES

**Table PO18. Induced Terminations of Pregnancy by Selected Characteristics and Age of Woman, New York City, 2015**

	Total	Age of Woman (Years)								Not Stated
		<18	18-19	20-24	25-29	30-34	35-39	≥40		
Induced Termination of Pregnancy, All	63,646	2,047	3,902	18,148	17,626	12,045	7,156	2,720	2	
Ethnic Group										
Hispanic	18,195	772	1,275	5,792	4,978	3,167	1,680	531	-	
Asian and Pacific Islander	4,028	47	135	836	1,118	933	640	319	-	
Non-Hispanic white	9,769	164	426	2,202	2,813	2,128	1,433	603	-	
Non-Hispanic black	25,698	902	1,712	7,588	7,051	4,762	2,682	1,000	1	
Other	2,155	65	157	693	595	339	234	72	-	
Unknown	3,801	97	197	1,037	1,071	716	487	195	1	
Marital Status										
Married	9,293	33	108	1,161	2,397	2,581	2,015	997	1	
Not married	46,402	1,828	3,398	15,043	12,998	7,785	4,060	1,290	-	
Other/Unknown	7,951	186	396	1,944	2,231	1,679	1,081	433	-	
Gestational Age (Weeks)										
≤ 6	24,560	565	1,257	6,642	7,332	4,838	2,828	1,098	-	
7 - 8	19,114	513	1,108	5,545	5,304	3,649	2,171	823	1	
9 - 10	8,354	332	577	2,457	2,222	1,510	949	307	-	
11 - 12	4,296	193	359	1,293	1,076	773	410	192	-	
13 - 15	3,035	165	210	861	760	546	360	132	1	
16 - 20	2,766	178	246	877	592	468	294	111	-	
≥ 21	1,485	101	143	459	331	255	141	55	-	
Unknown	36	-	2	14	8	6	3	2	-	
Type of Primary Termination Procedure										
Suction curettage	45,878	1,364	2,703	12,947	12,736	8,796	5,311	2,020	1	
Sharp curettage / D + C	1,382	45	83	306	331	292	203	121	1	
Dilatation and evacuation	5,284	305	437	1,563	1,226	935	589	229	-	
Intrauterine instillation	65	-	3	6	14	19	15	8	-	
Hysterotomy / hysterectomy	11	-	-	-	6	2	1	2	-	
Medical (non-surgical)	10,920	332	671	3,317	3,292	1,968	1,012	328	-	
Other	106	1	5	9	21	33	25	12	-	

Note: See Technical Notes: Spontaneous and Induced Terminations of Pregnancy.

New York City abortions by gestation 2015

## Second and third trimester abortion percentages by state, according to most recently published data

### North Carolina: 9.4 percent (residents only)

NC does not [break down](#) gestational age for all abortions reported -- just residents.

In [2016](#), there were 23,178 abortions reported. Of those, [2,184 abortions](#) out of 23,070 committed on North Carolina residents occurred at 13 weeks or greater: 1,398 (13-15 weeks), 785 (16-20 weeks), and 1 (21 weeks or greater). The percentage is likely greater when factoring in all abortions occurring in the state.

NC RESIDENT ABORTIONS: CHARACTERISTICS OF WOMEN RECEIVING ABORTIONS  
NORTH CAROLINA RESIDENTS, 2007 - 2016\*

	YEAR																			
	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016										
TOTALS	28,545	100.0	27,234	100.0	26,123	100.0	25,808	100.0	22,370	100.0	20,973	100.0	19,818	100.0	21,129	100.0	22,846	100.0	23,070	100.0
STATE OF OCCURRENCE:																				
... North Carolina	27,976	98.0	26,667	97.9	25,427	97.3	25,200	97.6	21,933	98.0	20,379	97.2	19,285	97.3	20,686	97.9	22,363	97.9	22,452	97.3
... Out of State	569	2.0	567	2.1	696	2.7	608	2.4	437	2.0	594	2.8	533	2.7	443	2.1	483	2.1	618	2.7
WEEKS OF GESTATION:																				
... 8 and under	17,043	59.7	17,074	62.7	16,079	61.6	16,632	64.4	14,540	65.0	12,875	61.4	12,131	61.2	13,040	61.7	14,060	61.5	15,153	65.7
... 9-12	6,479	22.7	6,102	22.4	5,726	21.9	5,631	21.8	4,859	21.7	4,369	20.8	4,188	21.1	4,233	20.0	4,487	19.6	4,934	21.4
... 13-15	1,739	6.1	1,715	6.3	1,408	5.4	1,445	5.6	1,252	5.6	1,175	5.6	1,016	5.1	1,049	5.0	1,305	5.7	1,398	6.1
... 16-20 (In-State Occurrences)	756	2.6	819	3.0	760	2.9	770	3.0	574	2.6	538	2.6	429	2.2	589	2.8	660	2.9	785	3.4
... 16-20 (Out-of-State Occurrences)	2	0.0	4	0.0	1	0.0	1	0.0	3	0.0	16	0.1	27	0.1	1	0.0	4	0.0	50	0.2
... 21+ (In-State Occurrences)	2	0.0	3	0.0	4	0.0	7	0.0	8	0.0	9	0.0	16	0.1	11	0.1	11	0.0	1	0.0
... 21+ (Out-of-State Occurrences)	1	0.0	6	0.0	2	0.0	0	0	0	0	68	0.3	39	0.2	1	0.0	5	0.0	35	0.2
... Unknown	2,523	8.8	1,511	5.5	2,143	8.2	1,322	5.1	1,134	5.1	1,923	9.2	1,972	10.0	2,205	10.4	2,314	10.1	714	3.1
TYPE OF PROCEDURE***																				
... Curettage	23,742	83.2	22,141	81.3	20,241	77.5	20,180	78.2	16,799	75.1	15,052	71.8	13,636	68.8	12,802	60.6	12,697	55.6	12,610	54.7
... Medical	3,645	12.8	4,275	15.7	4,730	18.1	4,920	19.1	5,236	23.4	5,127	24.4	5,061	25.5	7,287	34.5	9,267	40.6	9,748	42.3
... Intrauterine Instillation	183	0.6	140	0.5	176	0.7	37	0.1	10	0.0	5	0.0	10	0.1	5	0.0	3	0.0	1	0.0
... Hysterectomy/Hysterotomy	0	0	2	0.0	0	0	0	0	0	0	1	0.0	0	0	0	0	0	0	0	0
... Unknown	975	3.4	676	2.5	976	3.7	671	2.6	325	1.5	788	3.8	1,111	5.6	1,035	4.9	879	3.8	711	3.1

North Carolina abortions for residents by gestation 2007 to 2016

### North Dakota: 9.3 percent

In [2016](#), 114 out of 1,226 abortions were committed on or after the 13th week: 34 (13 weeks), 32 (14 weeks), 23 (15 weeks), and 25 (16 weeks or greater).

ND OCCURRENCE INDUCED TERMINATION OF PREGNANCY  
2016

#### INDUCED TERMINATION OF PREGNANCY BY WEEKS GESTATION

WEEKS GESTATION	NUMBER OF EVENTS
UNKNOWN	0
FIVE OR LESS	154
SIX	251
SEVEN	213
EIGHT	183
NINE	133
TEN	82
ELEVEN	54
TWELVE	42
THIRTEEN	34
FOURTEEN	32
FIFTEEN	23
SIXTEEN OR MORE	25

OCCURRENCE TOTAL 1226

North Dakota abortions by gestation 2016



## Second and third trimester abortion percentages by state, according to most recently published data

### Ohio: 15.4 percent

In [2016](#), 3,192 out of 20,672 abortions were reported on or after 13 weeks: 2,684 (13-18 weeks), 368 (19-20 weeks), and 140 (21 weeks and up).

Table 2. Selected Characteristics of Induced Abortions Reported in Ohio, 2006-2016

	CHARACTERISTIC	2016	2015	2014	2013	2012	2011	2010	2009	2008	2007	2006
NUMBER OF ABORTIONS												
	TOTAL INDUCED ABORTIONS	20,672	20,976	21,186	23,216	25,473	24,764	28,123	28,721	29,613	30,859	32,936
Residence	Ohio Resident	19,543	19,765	20,018	22,011	24,080	23,250	26,322	26,959	27,672	28,921	30,809
	Out-of-State Resident	1,129	1,211	1,168	1,205	1,393	1,511	1,801	1,762	1,941	1,938	2,127
Completed Weeks of Gestation	Less than 9 Weeks	11,230	10,910	11,088	13,128	14,364	14,105	16,283	16,264	16,663	17,023	17,955
	9-12 Weeks	6,250	6,632	6,624	6,624	7,220	6,909	7,672	7,971	8,257	8,855	9,862
	13 - 18 Weeks	2,684	2,956	2,964	2,925	3,176	2,897	3,223	3,390	3,629	3,589	3,699
	19-20 Weeks	368	333	377	359	445	318	345	340	339	410	448
	21 Weeks and Over	140	145	133	173	180	378	458	480	480	502	510
	Not Reported	0	0	0	7	88	157	142	276	245	480	462

Ohio abortions by gestation 2006-2016

### Oklahoma: 7 percent

In [2017](#), 327 abortions out of 4,723 were performed between 13-21 weeks as follows: 174 (13-15 weeks), 132 (16-20 weeks), and 21 (21 weeks).

**Requirement 17:** The number of abortions resulting in reported complications, and of those, how many were reported by the physician who performed the abortion, and how many were reported by another physician, the types of reported complications, and the number of each type based on data which shall be compiled and transmitted to the State Department of Health by the State Board of Medical Licensure and Supervision and the State Board of Osteopathic Examiners.

Table 24. Complications Reported by

The number of abortions resulting in complications:	Frequency	Percent
Reported by physician who performed abortion	29	93.6
Another physician	**	
<b>Types of Complications*</b>		
Failed termination of pregnancy (continued viable pregnancy)	22	71.0
Heavy bleeding/hemorrhage with estimated blood loss of greater than or equal to 500cc	**	**
Other	7	22.6

\*More than one complication can be reported for the same event

\*\*Cell is suppressed to maintain confidentiality of surrounding entries

Abortion complications reported Oklahoma 2017

### Oregon: 11 percent

In [2016](#), 965 abortions out of 8,942 were reported on or after the 13th week: 537 (13-16 weeks), 254 (17-20 weeks), 96 (21-22 weeks) and 78 (23 weeks and up). Preliminary stats reveal that 8,107 abortions have been reported in [2017](#), however, further data has not been released as of the date of this report.

Induced termination of pregnancy

3-1

TABLE 3-4. Abortions in relation to length of gestation by method, complications, and age of patient, Oregon occurrence, 2016

Method, complications and age of patient	Total	Weeks gestation						
		< 9	9-12	13-16	17-20	21-22	23+	Unk.
<b>Total</b>	8,942	6,452	1,501	537	254	96	78	24
<b>Method</b>								
Suction curette	3,043	1,826	992	213	3	1	—	8
Medical (non-surgical)	3,880	3,670	166	1	7	9	14	13
Dilation & evacuation	1,999	952	343	322	242	81	57	2
Vaginal prostaglandin	11	—	—	1	2	3	5	—
Sharp curettage	1	1	—	—	—	—	—	—
Other	8	3	—	—	—	2	2	1

2016 Oregon Abortion by gestation



## Second and third trimester abortion percentages by state, according to most recently published data

### Pennsylvania: 13 percent

In [2016](#), 3,910 abortions out of 30,881 were reported between 13-24 weeks: 1,549 (13-14 weeks), 1,251 (15-17 weeks), 719 (18-20 weeks), 391 (23 weeks), and zero (24 weeks and up).

Pennsylvania Induced Abortion Report  
January - December 2016

Table 1 Induced Abortions Performed In Pennsylvania by Age of Woman		Table 2 Induced Abortions Performed In Pennsylvania by Weeks of Gestation	
Age (Years)	Number	Weeks of Gestation	Number
12 or Under	1	8 or Less	19,657
13	19	9 - 10	4,878
14	57	11 - 12	2,436
15	146	13 - 14	1,549
16	272	15 - 17	1,251
17	393	18 - 20	719
18	790	21 - 23	391
19	1,127	24 or More	0
20 - 24	9,502	Total	30,881
25 - 29	9,211		
30 - 34	5,428		
35 - 39	2,941		
40 - 44	918		
45 or Older	76		
Unknown	0		
Total	30,881		

Note: Weeks of gestation are based on clinical estimate.

Pennsylvania abortions by gestation 2016

### Rhode Island: unknown

### South Carolina: <1 percent, most likely higher

This state does not break down abortion stats in the 13th week. In [2017](#), 25 out of 3,875 abortions were reported from 14-23 weeks: 19 (14-19 weeks) and 6 (20-23 weeks). However, 3,222 abortions were committed from 7-13 weeks, so it is likely that this percentage would be much higher, if the numbers in the 13th week were specified.

Table 3A.  
Abortions by Method and Clinical Estimate of Gestation  
South Carolina, 2017 (Jan 1 - Sept 24)

Method of Abortion	Total	Weeks Gestation					
		Under 6	7 - 13	14 - 19	20 - 23	24 or more	Unknown
Total	3,875	1,321	2,531	17	6	-	-
Suction Curettage	1,623	251	1,370	2	-	-	-
Dilation and Evacuation	11	-	-	10	1	-	-
Intra-Uterine Saline Instillation	1	-	-	-	1	-	-
Intra-Uterine Prostaglandin Instillation	3	-	-	2	1	-	-
Hysterotomy/Hysterectomy	1	-	-	1	-	-	-
Medical	2,235	1,070	1,161	1	3	-	-
Other	1	-	-	1	-	-	-

Table 3B.  
Abortions by Method and Probable Postfertilization Age  
South Carolina, 2017 (Sept 25 - Dec 31)


Method of Abortion	Total	Weeks Gestation					
		Under 6	7 - 13	14 - 19	20 - 23	24 or more	Unknown
Total	1,237	544	691	2	-	-	-
Dilation and Curettage	128	22	106	-	-	-	-
Manual Vacuum Aspiration	89	39	50	-	-	-	-
Electrical Vacuum Aspiration	300	66	234	-	-	-	-
Dilation and Evacuation	5	-	4	1	-	-	-
Combined Induction Abortion and Dilation and Evacuation	-	-	-	-	-	-	-
Medication Abortion	714	417	296	1	-	-	-
Induction Abortion with Prostaglandins	1	-	1	-	-	-	-
Induction Abortion with Intra-Amniotic Instillation	-	-	-	-	-	-	-
Induction Abortion - other	-	-	-	-	-	-	-
Hysterotomy / Hysterectomy	-	-	-	-	-	-	-
Intact Dilation and Extraction (partial birth)	-	-	-	-	-	-	-
Other	-	-	-	-	-	-	-

SC abortions 2017

## Second and third trimester abortion percentages by state, according to most recently published data

### South Dakota: 6 percent

In [2016](#), 26 out of 472 abortions were committed at 13 weeks or greater.



South Dakota abortions by weeks 2016

### Tennessee: 11.2 percent

In [2016](#), 1,092 out of 9,732 abortions were committed at 13 weeks or greater: 788 (13-14 weeks), 207 (15-16 weeks), and 97 (17-20 weeks). There were 116 abortions reported at unknown gestation.

#### SELECTED INDUCED TERMINATION OF PREGNANCY (ITOP) DATA, ACCORDING TO AGE AND RACE OF WOMAN, TENNESSEE AND DEPARTMENT OF HEALTH REGIONS, RESIDENT DATA, 2016

TENNESSEE			NUMBER OF PREVIOUS ITOPs			GESTATIONAL AGE		
MARITAL STATUS			NUMBER	FREQ	PCNT	WEEKS	NUMBER	PCNT
MARRIED	1,239	12.7	0	6,096	62.6	LT. 7	2,143	22.0
NOT MARRIED	8,287	85.2	1	2,317	23.8	7-8	3,356	34.5
UNKNOWN	206	2.1	2	793	8.1	9-10	1,909	19.6
TOTAL	9,732	100.0	3	268	2.8	11-12	1,116	11.5
			4+	153	1.6	13-14	788	8.1
			UNKNOWN	105	1.1	15-16	207	2.1
			TOTAL	9,732	100.0	17-20	97	1.0
						UNKNOWN	116	1.2
						TOTAL	9,732	100.0

Tennessee abortions by gestation 2016

### Texas: 6.2 percent

In [2015](#), 3,431 out of 55,287 were [committed](#) at 13 weeks or greater including 25 in the third trimester: 1,556 (13-14 weeks), 913 (15-16 weeks), 867 (17-20 weeks), 69 (21-24 weeks), and 26 (25 weeks or greater).

Table 36 Induced Terminations of Pregnancy by Procedure and Length of Gestation Texas

TABLE 36  
INDUCED TERMINATIONS OF PREGNANCY BY PROCEDURE  
AND POST-FERTILIZATION AGE - 2015\*

PROCEDURE	TOTAL	< 9 WEEKS	9 - 10 WEEKS	11 - 12 WEEKS	13 - 14 WEEKS	15 - 16 WEEKS	17 - 20 WEEKS	21 - 24 WEEKS	>= 25 WEEKS	UNK
SUCTION CURETTAGE	43,823	36,834	4,235	2,149	464	122	15	1	0	3
MEDICAL (Non Surgical)	6,029	5,971	3	3	2	6	31	10	2	1
DILATION & EVACUATION	4,386	818	344	664	1,065	764	728	3	0	0
INTRA-UTERINE	3	1	0	1	0	1	0	0	0	0
SHARP CURETTAGE	68	56	1	6	4	0	1	0	0	0
HYSTEROTOMY/HYSTERECTOMY	4	0	0	0	0	0	4	0	0	0
OTHER/ UNKNOWN	974 **	679	40	27	21	20	88	55	24	20
TOTAL	55,287	44,359	4,623	2,850	1,556	913	867	69	26	24

Texas abortions by gestation 2015

## Second and third trimester abortion percentages by state, according to most recently published data

### Utah: at least 9 percent (residents only)

In [2016](#), 3,008 abortions were reported in the state but gestational age was only reported for residents. Out of the 2,808 abortions for residents, 248 were committed on or after the 13 week as follows: 111 (13-14 weeks), 120 (15-20 weeks), and 17 (21 weeks and greater). This percentage is likely higher if it included all abortions occurring in the state.

**Table R12. Abortions by weeks gestation and reason for termination, residents: Utah, 2016**

Weeks gestation	Total	Reason for termination				
		Maternal life endangered	Therapeutic	Fetal malformation	HIV positive	Elective
<b>Total</b>	2,808	15	2,515	41	**	207
<b>Under 5</b>	6	**	6	**	**	**
<b>5-6</b>	1,040	**	954	**	**	62
<b>7-8</b>	901	**	819	**	**	70
<b>9-10</b>	351	**	315	**	**	31
<b>11-12</b>	234	**	201	**	**	30
<b>13-14</b>	111	**	92	5	**	10
<b>15-20</b>	120	**	95	22	**	3
<b>21 and over</b>	17	**	7	9	**	0

The last menstrual date is used to calculate gestation when the physician's clinical estimate is not available.

Numbers may not add to total due to missing data.

\*\*Small counts (<5) are not displayed.

Utah abortions by gestation 2016 residents by reason

### Vermont: 8.3 percent

In [2016](#), 108 out of 1,298 abortions were committed at 13 weeks or greater as follows: 63 (13-15 weeks), 31 (16-20 weeks), and 14 (21 weeks or greater).

**TABLE E-4  
2016 VERMONT ABORTIONS - OCCURRENCE**

WEEKS GESTATION BY AGE OF PATIENT								
	WEEKS GESTATION <sup>(1)</sup>							
AGE OF PATIENT	< 9	9 - 10	11 - 12	13 - 15	16 - 20	21 +	UNKNOWN	TOTAL
< 15 YEARS	0	0	0	1	0	0	0	1
15 YEARS	3	2	1	0	0	0	0	6
16 YEARS	8	3	2	0	1	0	0	14
17 YEARS	17	7	0	2	1	0	0	27
18 YEARS	28	10	4	4	1	0	2	49
19 YEARS	42	9	2	1	1	1	0	56
20-24 YEARS	256	71	25	21	3	1	0	377
25-29 YEARS	259	50	23	19	8	2	1	362
30-34 YEARS	170	28	17	8	7	6	0	236
35-39 YEARS	76	14	5	4	5	2	0	106
40+ YEARS	33	6	8	3	3	2	0	55
UNKNOWN	5	1	2	0	1	0	0	9
TOTAL	897	201	89	63	31	14	3	1298

<sup>(1)</sup> SEE APPENDIX B FOR DATA QUALITY NOTES.

Vermont abortions by gestation 2016

### Virginia: unknown

Although Virginia [publishes](#) data on abortions, they do not publish data on how many occurred by week of gestation.

## Second and third trimester abortion percentages by state, according to most recently published data

### Washington State: 11.4 percent

In [2016](#), 1,950 abortions out of 17,093 were committed at 13 weeks or greater: 474 (13-15 weeks), 980 (16-19 weeks), and 496 (20 weeks and over).

A. Age of Woman			B. Weeks of Gestation <sup>2</sup>		
Age	Number	Percent <sup>1</sup>	Weeks of Gestation	Number	Percent <sup>1</sup>
State Total	17,093	100.0	State Total	17,093	100.0
Under 15	28	0.2	1-4	9	0.1
15-19	1,786	10.4	5-8	460	2.7
15-17	567	3.3	9	11,693	68.4
18-19	1,219	7.1	10	1,283	7.5
20-24	4,856	28.4	11	699	4.1
25-29	4,715	27.6	12	623	3.6
30-34	3,203	18.7	13-15	474	2.8
35-39	1,798	10.5	16-19	980	5.7
40-44	634	3.7	20 and Over	496	2.9
45 and Over	56	0.3	Unknown	348	2.0
Unknown	17	0.1			

Washington State abortions by gestation 2016

### West Virginia: 9 percent

In [2015](#), 136 out of 1,519 abortions were committed at 13 weeks or greater as follows: 93 (13-15 weeks), 38 (16-19 weeks), and 5 (20 weeks or greater). The report indicates that there were four abortions committed due to a “medical emergency,” but does not state what the emergency was.

Induced Terminations of Pregnancy  
By Age of Female and Weeks of Gestation  
West Virginia Occurrences, 2015  
Table 33

Age of Female	Weeks of Gestation							Total
	Less than 9	9-10	11-12	13-15	16-19	20 or More	Unknown	
10-14	3	0	1	1	0	1	0	6
15-17	21	8	10	3	0	0	0	42
18-19	65	20	12	4	2	0	0	103
20-24	294	83	60	28	14	1	0	480
25-29	232	77	48	23	13	1	0	394
30-34	182	49	44	23	6	1	0	305
35-39	98	21	14	10	1	1	0	145
40-44	21	10	5	1	1	0	0	38
45+	2	0	0	0	0	0	0	2
Unknown	1	1	1	0	1	0	0	4
Total	919	269	195	93	38	5	0	1,519

Note: Gestational age is the physician's estimate of gestation (clinical/obstetric estimate).

If the physician's estimate of gestation is not available, it is calculated from the date of onset of last normal menses.

West Virginia abortions by gestation 2015

## Second and third trimester abortion percentages by state, according to most recently published data

### Wisconsin: 15 percent

In [2016](#), 835 out of 5,612 were committed at 13 weeks or greater: 445 (13-15 weeks) and 353 (16 – 20 weeks), and 37 (over 20 weeks).

Table 8. Reported Induced Abortions by Estimated Gestation, Wisconsin 2016

Weeks of Gestation	Occurred in Wisconsin		Wisconsin Residents		Percent Wisconsin Residents
	Number	Percent	Number	Percent	
8 weeks or less	3,116	56	3,048	56	98
9-10	1,113	20	1,080	20	97
11-12	548	10	534	10	97
13-15	445	8	430	8	97
16-20	353	6	343	6	97
Over 20	37	1	37	1	100
Total	5,612	100	5,472	100	98

Source: Office of Health Informatics, Division of Public Health, Wisconsin Department of Health Services.

Note: The "Wisconsin Residents" columns include only induced abortions that were obtained in Wisconsin by Wisconsin residents. Information about Wisconsin residents who obtained induced abortions in other states is not available.

The data provided for what is termed "weeks of gestation" in the tables cannot be classified precisely or consistently. In some cases, it refers to the patient's reported weeks since last menstrual period; in other cases, it is the physician's estimate of weeks since conception. Reporting instructions ask for the number of weeks since patient's last menstrual period. If this information is unavailable, reporters are asked to substitute a clinical estimate of gestation. It is not known which reports used which measure, nor the proportion of reports using each. A physician's estimate of gestational age based on clinical observation is generally more reliable and may be several weeks less than the patient's reported weeks since last menstrual period. Readers should be aware of the imprecision in the measure. The table groups the reported weeks into categories used by the U.S. Centers for Disease Control and Prevention and the Alan Guttmacher Institute.

Wisconsin abortions by gestation 2016

### Wyoming: unknown

For the sake of consistency, unless otherwise noted, the percentages used were based on gestational age or estimated gestational age. Live Action News did not locate gestational data for fourteen states and the District of Columbia.

*Updated: July 10, 2018*